

# Balloon in the Flask

## What is Pressure?

### Introduction

Heat some water in a flask, then attach a balloon, cool the flask, and watch as the balloon collapses into the flask. An easy-to-perform and colorful variation of the common *Crush the Can* demonstration.

### Concepts

- Pressure differential
- Vacuum

### Materials

- Erlenmeyer flask, borosilicate glass, 250-mL
- Balloon, latex, 11-inch size (size to fit flask)
- Hot plate or Bunsen burner
- Ice bath or cold running water
- Water, 25 mL



### Safety Precautions

Always practice a demonstration before presenting it to students. Be careful of the hot glass and steam. Wear chemical splash goggles and heat-resistant gloves.

### Procedure

1. Add approximately 25 mL of water to a 250-mL Erlenmeyer flask. Heat the water using a hot plate, Bunsen burner, or other heat source.
2. As the water comes to a boil and steam begins to rise out of the flask, remove the flask from the heat. Quickly place the balloon over the mouth of the flask.
3. Place the flask under cold running water and the balloon will be pushed into the flask until it fills the entire flask. If the balloon stretches too much, it may break.



### Tips

- Use a borosilicate (Pyrex®) flask with a heavy-duty rim. Do not use an economy-choice flask. Check the flask for chips or cracks before use.
- Stretch out the balloon by inflating and deflating it before using it.
- The demonstration works best if the balloon is centered on the opening when placed over the mouth of the flask. It also helps if the balloon is slightly pushed into the flask when it begins to collapse. If not, it may collapse onto itself and not get drawn into the flask. The demonstration will work without holding it under cold water, but it takes longer to cool the glass and condense the water vapor.
- A hard-boiled, shelled egg can also be used in place of the balloon. A larger flask may be needed depending on the size of the egg.

## Discussion

The *Balloon in the Flask* demonstration is an easy-to-perform and colorful variation of the common *Crush the Can* demonstration. Both demonstrations rely on the creation of a pressure differential caused by the condensation of steam inside a closed system. As the steam condenses, a partial vacuum is formed inside the closed system. The pressure outside the closed system is still at atmospheric pressure (approximately 14.7 lb/in<sup>2</sup>). This pressure difference will cause the balloon to be pushed into the flask. The balloon is not “sucked” into the flask—it is pushed in by the greater atmospheric pressure that exists outside the closed system. The balloon will continue to be pushed into the flask until the pressure inside the closed system is approximately equal to the atmospheric pressure.

## Connecting to the National Standards

This laboratory activity relates to the following National Science Education Standards (1996):

### *Unifying Concepts and Processes: Grades K–12*

Systems, order, and organization

Evidence, models, and explanation

### *Content Standards: Grades 5–8*

Content Standard B: Physical Science, properties and changes of properties in matter, understanding of motions and forces

### *Content Standards: Grades 9–12*

Content Standard B: Physical Science, structure and properties of matter, motions and forces

## Reference

Shakhashiri, B. Z. *Chemical Demonstrations: A Handbook for Teachers in Chemistry*; University of Wisconsin: Madison; Vol. 2, pp 6–8.

## Flinn Scientific—Teaching Chemistry™ eLearning Video Series

A video of the *Balloon in the Flask* activity, presented by George Gross, is available in *What Is Pressure?*, part of the Flinn Scientific—Teaching Chemistry eLearning Video Series.

## Materials for *Balloon in the Flask* are available from Flinn Scientific, Inc.

Catalog No.	Description
AP1900	Balloons, Latex, pkg/20
AP8386	Hot Plate, Flinn, 7" × 7"
GP3045	Erlenmeyer Flask, Borosilicate Glass, 250-mL

Consult your *Flinn Scientific Catalog/Reference Manual* for current prices.